

Primitive Forms, Topological LG Models Coupled to Gravity and Mirror Symmetry

Atsushi Takahashi *

Research Institute for Mathematical Sciences,
Kyoto University,
Kyoto 606-01, Japan

Abstract

In this paper, we will describe the mathematical foundation of topological Landau–Ginzburg (LG) models coupled to gravity at genus 0 in terms of primitive forms. In particular, we can give a natural explanation for the large phase space.

We also discuss the mirror symmetry for Calabi–Yau manifolds and $\mathbb{CP}^1$ in our context. We see that the mirror partner of $\mathbb{CP}^1$ is the theory of primitive form associated to $f = z + qz^{-1}$.

Introduction

Mirror symmetry (for detail see [GY][Y]) is the one of the most important phenomena in algebraic geometry and string theory. A pair of manifolds (X, Y) is said to be mirror pair if A-twisted sigma model coupled to gravity on X is isomorphic to B-twisted sigma model coupled to gravity on Y [W3]. The correlation functions of A-twisted sigma model coupled to gravity have been mathematically established as the theory of Gromov–Witten invariants [Beh][BF][BM][KM1][LT1][LT2][LT3][RT1][RT2] and have been intensively investigated by several authors [EHX1][EHX2][EX][G1][G2][KM2].

* *E-mail address* : atsushi@kurims.kyoto-u.ac.jp

On the other hand, B-twisted sigma models coupled to gravity has not been understood mathematically (at least for $g > 0$ sector) but topological Landau–Ginzburg models coupled to gravity have been studied based on singularity theory [LVW][VW][Mar]. Recently, Barannikov and Kontsevich have proposed a new construction of Frobenius manifolds for Calabi–Yau manifolds with arbitrary dimension [BK], which is conjectured to be a mirror partner for that of Gromov–Witten theory.

The purpose of this paper is to describe the mathematical foundation of topological Landau–Ginzburg models coupled to gravity at genus 0 in terms of the theory of primitive forms which was introduced by K. Saito [S1][S2] in his study for the period mapping for a universal unfolding of a function with an isolated critical singularity. Then, we discuss the mirror symmetry for Calabi–Yau manifolds and $\mathbb{CP}^1$. Especially, we can show that the mirror partner of $\mathbb{CP}^1$ is the theory of primitive forms associated to a function $f = z + qz^{-1}$ on $\mathbb{C}^*$. This reproduce the results in [EHX2].

In section 1, we review the theory of primitive forms [S1][S2]. Some definitions and notations are modified so that they can be identified with those in physics literatures. We will first introduce the notion of a Landau–Ginzburg system $F(z, t)$ over a frame (Z, X, S, T) , which corresponds to perturbation theory of the original Landau–Ginzburg models. Then we will introduce the chiral ring structure (the residual product $\circ$), the perturbed large phase space $\pi_*\mathcal{H}_F^{(0)}$, the Gauss–Manin connection ∇ on $\pi_*\mathcal{H}_F^{(0)}$, the Euler vector field E and the higher residue pairings $K^{(k)}$. We will see that a primitive form $\zeta^{(0)}$ associated to F can be defined if the above data are given.

In section 2, we first review the Frobenius structure defined in [D] [Man1]. We will show that the theorem.

Theorem. *Let $F(z, t)$ be an LG system over a frame (Z, X, S, T) and let us assume that a primitive form $\zeta^{(0)}$ is given. Then S is a Frobenius manifold with an identity δ_0 and an Euler vector field $E = w(\delta_0)$.*

We will construct Cohomological Field Theories [KM1][Man1] associated to primitive forms. We will see that the large phase space and the gravitational descendants are given by $H_F = \pi_*\mathcal{H}_F^{(0)}/m_T\pi_*\mathcal{H}_F^{(0)}$ and $\nabla_{\delta_i}\zeta^{(-d-1)}|_{t'=0}$, respectively. Then we see that

Theorem. *Let $F(z, t)$ be an LG system over a frame (Z, X, S, T) . Then a primitive form $\zeta^{(0)}$ uniquely defines the genus 0 free energy Φ_F^{grav} up to quadratic terms, which is a generating function of correlation functions of the LG models coupled gravity whose superpotential is given by $f(z) = F(z, 0)$.*

Note that if we want to have a Frobenius structure, we must fix a primitive form $\zeta^{(0)}$ for the LG system F .

In section 3, we discuss the relations between the primitive forms and the mirror symmetry. We will first review the period mapping of the primitive forms. Then we will show that the canonical coordinates with integral exponents have a geometrical meaning, i.e., which are one to one correspondence with the primitive part of the cohomology of V . In particular, if V is a smooth Calabi–Yau manifold, we can consider that the Frobenius submanifold S' of S (integral exponents sector) is a Frobenius submanifold of that of Barannikov–Kontsevich construction from the viewpoint of the LG orbifold theory [IV]. We will also show that a relation between primitive forms and (normalized) holomorphic forms on Calabi–Yau manifolds. Finally, we will describe the mirror symmetry of $\mathbb{CP}^1$ in terms of primitive forms;

Theorem. *A primitive form for LG system*

$$F(z, t) = t^0 + z + q \exp(t^1) z^{-1} \quad (1)$$

is given by

$$\zeta^{(0)} = \frac{dz}{z}. \quad (2)$$

Then we have

$$\Phi_{\mathbb{CP}^1}^{st}(t) = \Phi_F^{grav}(\tilde{t}), \quad (3)$$

where $\Phi_{\mathbb{CP}^1}^{st}$ is the (large phase space) stable genus 0 generating function for Gromov–Witten Invariants

$$\Phi_{\mathbb{CP}^1}^{st}(t) := \sum_{n \geq 3, (a_i, d_i)} \frac{1}{n!} t_{d_1}^{a_1} \cdots t_{d_n}^{a_n} \sum_{\beta \in \mathbb{Z}_{\geq 0}} q^\beta \langle \sigma_{d_1}(\mathcal{O}_{a_1}) \cdots \sigma_{d_n}(\mathcal{O}_{a_n}) \rangle_{0, \beta[\mathbb{CP}^1]} \quad (4)$$

and

$$\tilde{t}_d^i := t_d^i + \sum_{(j, e)} t_e^j \sum_{\beta \in \mathbb{Z}_{\geq 0}} q^\beta \langle \sigma_{e-d-1}(\mathcal{O}_j) \mathcal{O}^i \rangle_{0, \beta[\mathbb{CP}^1]}. \quad (5)$$

This shows that the pair $(\mathbb{C}^*, f)$ is a mirror of $\mathbb{CP}^1$.

This approach may be valid for the other toric Fano manifolds or Grassmannian manifolds whose quantum cohomology rings are given by the Jacobian ring [Bat][EHX2].

We can apply the notion of primitive forms to the Seiberg–Witten theory. For example, the Seiberg–Witten form λ_{SW} for A_1 is a meromorphic 1-form on

$$F(x, t^0, \mu) = W_{A_1}(x, t^0 + z + \mu^2 z^{-1}) = x^2 + t^0 + z + \mu^2 z^{-1} = 0,$$

and given by $\lambda_{SW} = x \frac{dz}{z}$. In this case, the Seiberg–Witten form exactly coincides with the primitive form $(\zeta^{(-1)})$ for the above defining equation $F(x, t^0, \Lambda)$. For other gauge groups, we see that λ_{SW} is given by the “joint” of the primitive forms for $\mathbb{C}^*$ and ADE -type singularity and reproduces the results of [IY]. We will discuss this in detail in the next paper.

After the completion of this paper, we noticed a preprint by Manin [Man2] where the construction of Frobenius manifolds by primitive forms is presented.

Acknowledgement

I am deeply grateful to Professor Kyoji Saito for his encouragement. I also would like to thank Professor Tohru Eguchi, Professor Shinobu Hosono and Professor Toshiya Kawai for useful discussions.

Contents

1	LG system and primitive form	5
1.1	Landau–Ginzburg system	5
1.2	The residual product $\circ$	6
1.3	De Rham cohomology $\mathcal{H}_F^{(0)}$	8
1.4	Higher residue pairings $K^{(k)}$ and primitive form $\zeta^{(0)}$	10
1.5	Canonical coordinate system	11
2	Cohomological Field Theory associated to LG system	13
2.1	Frobenius structure	13
2.2	Cohomological Field Theory	15
3	Primitive Forms and Mirror Symmetry	18
3.1	Period of Primitive Forms	18
3.2	Landau–Ginzburg Orbifolds and Calabi–Yau Varieties	21
3.3	Mirror Symmetry for $\mathbb{CP}^1$	23

1 LG system and primitive form

1.1 Landau–Ginzburg system

Let us first take the holomorphic function f (*superpotential*),

$$f(z) : (\mathbb{C}^{n+1}, 0) \rightarrow (\mathbb{C}, 0),$$

which is a quasi-homogeneous function of z_i and has an isolated singularity at base point 0. Then the *Jacobian ring* (or *chiral ring*) $\mathcal{R}_0$ defined by

$$\mathcal{R}_0 := \mathcal{O}_{\mathbb{C}^{n+1}, 0} \Big/ \left(\frac{\partial f}{\partial z_0}, \dots, \frac{\partial f}{\partial z_n} \right)$$

becomes a finite dimensional $\mathbb{C}$ -vector space and we denote $\dim_{\mathbb{C}} \mathcal{R}_0$ by μ .

Let us consider a pair $(S, \delta_0, 0)$ where S is a μ -dimensional complex manifold with a base point $0 \in S$ and δ_0 is a holomorphic vector field on S which is non-singular at 0.

There exist a $(\mu - 1)$ -dimensional manifold T with a base point $0 \in T$ and a holomorphic submersion π defined locally,

$$\pi : (S, 0) \rightarrow (T, 0) \text{ such that } \pi^{-1}\mathcal{O}_T = \{g \in \mathcal{O}_S : \delta_0 g = 0\} \quad (6)$$

where $\mathcal{O}_S$ and $\mathcal{O}_T$ are structure sheaves of S and T . By choosing neighborhoods of base points of S and T , we may assume that T is the orbit space of δ_0 . Let us denote by Der_S the sheaf of germs of holomorphic vector fields on S .

Put

$$\mathcal{G} := \{\delta \in \pi_* Der_S : [\delta_0, \delta] = 0\} \quad (7)$$

where $[\ , \]$ is the bracket product of vector fields.

The module $\mathcal{G}$ is naturally $\mathcal{O}_T$ -free module of rank μ closed under the Lie-bracket.

Remark. We have the following exact sequence of $\mathcal{O}_T$ -Lie algebras:

$$0 \rightarrow \mathcal{O}_T \delta_0 \rightarrow \mathcal{G} \rightarrow Der_T \rightarrow 0. \quad (8)$$

In this situation, the following are equivalent.

1. To give a holomorphic function t^0 on $(S, \delta_0, 0)$ such that $t^0(0) = 0$ and $\delta_0 t^0 = 1$.
2. To give a splitting of the above exact sequence as an $\mathcal{O}_T$ -Lie algebra.

Let X be an $(n + \mu)$ -dimensional complex manifold with a base point $0 \in X$, and let $q : (X, 0) \rightarrow (T, 0)$ be a smooth surjective map. Let $(Z, 0)$ be the fiber product with the diagram,

$$\begin{array}{ccc} (Z, 0) & \xrightarrow{\hat{\pi}} & (X, 0) \\ p \downarrow & & \downarrow q \\ (S, 0) & \xrightarrow{\pi} & (T, 0) \end{array} \quad (9)$$

There exists a unique lifting $\hat{\delta}_0$ on Z of the primitive vector field δ_0 on S such that $\hat{\pi} : (Z, 0) \rightarrow (X, 0)$ is the projection to the orbit space of $\hat{\delta}_0$.

We shall call the above diagram a **frame**, which will often be denoted simply by (Z, X, S, T) .

Remark. The following are equivalent.

1. To give a holomorphic function $F(z, t)$ on Z such that $\hat{\delta}_0 F = 1$ and $F(0) = 0$.
2. To give a holomorphic mapping $\varphi : (X, 0) \rightarrow (S, 0)$, such that $\pi \circ \varphi = q$.

Definition 1.1. A holomorphic function $F(z, t)$ on Z is called a *Landau–Ginzburg system (LG system)* over a frame (Z, X, S, T) if it satisfies the following three conditions:

1. $F(z, 0) = f(z)$.
2. $\hat{\delta}_0 F = 1$, $F(0) = 0$.
3. The correspondence

$$T(S)_0 \rightarrow \mathcal{R}_0 = \mathcal{O}_{p^{-1}(0), 0} \left/ \left(\frac{\partial f}{\partial z_0}, \dots, \frac{\partial f}{\partial z_n} \right) \right., \quad \delta \mapsto \hat{\delta}F|_{p^{-1}(0)}, \quad (10)$$

is bijective.

1.2 The residual product $\circ$

Let $F(z, t)$ be an LG system over a frame (Z, X, S, T) and let $\varphi : (X, 0) \rightarrow (S, 0)$ be an associated holomorphic map.

Let $\mathcal{C}$ be the subvariety of the critical points of φ in X defined by the ideal $(\frac{\partial F}{\partial z_0}, \dots, \frac{\partial F}{\partial z_n})\mathcal{O}_X$. Then we see that the restriction $q|_{\mathcal{C}} : \mathcal{C} \rightarrow T$ is a μ -sheeted branched covering and $q_*\mathcal{O}_{\mathcal{C}}$ is an $\mathcal{O}_T$ -free module of rank μ .

Let us put $\hat{\mathcal{C}} := \hat{\pi}^{-1}\mathcal{C} \subset Z$, which is defined by the ideal $(\frac{\partial F}{\partial z_0}, \dots, \frac{\partial F}{\partial z_n})\mathcal{O}_Z$. Viewing X as a hypersurface in Z defined by $F = 0$, we have the following exact sequence:

$$0 \rightarrow p_*\mathcal{O}_{\hat{\mathcal{C}}} \xrightarrow{p_*F} p_*\mathcal{O}_{\hat{\mathcal{C}}} \rightarrow \varphi_*\mathcal{O}_{\mathcal{C}} \rightarrow 0, \quad (11)$$

where p_*F means an operator of a multiplication of F .

Lemma 1.1. *Let $F(z, t)$ be an LG system over a frame (Z, X, S, T) . Then the $\mathcal{O}_S$ -homomorphism*

$$Der_S \rightarrow p_*\mathcal{O}_{\hat{\mathcal{C}}}, \quad \delta \mapsto \hat{\delta}F|_{\hat{\mathcal{C}}}, \quad (12)$$

and the $\mathcal{O}_T$ -homomorphism

$$\mathcal{G} \rightarrow q_*\mathcal{O}_{\mathcal{C}}, \quad \delta \mapsto \hat{\delta}F|_{\mathcal{C}}. \quad (13)$$

are bijections.

Corollary. *We can give a ring structure and a t^0 -multiplication structure on $\mathcal{G}$ by,*

$$(\delta \hat{\circ} \delta')F|_{\mathcal{C}} := \hat{\delta}F|_{\mathcal{C}} \cdot \hat{\delta}'F|_{\mathcal{C}} \quad (14)$$

and

$$(t^0 \hat{\circ} \delta)F|_{\mathcal{C}} := t^0|_{\mathcal{C}} \cdot \hat{\delta}F|_{\mathcal{C}}. \quad (15)$$

Definition 1.2. We call the above $\circ$ the *residual product*.

Since $p|_{\hat{\mathcal{C}}} : \hat{\mathcal{C}} \rightarrow S$ is μ -sheeted branched covering, $p_*\mathcal{O}_{\hat{\mathcal{C}}}$ is an $\mathcal{O}_S$ -free module of rank μ , and hence the sequence is an $\mathcal{O}_S$ -free resolution of $\varphi_*\mathcal{O}_{\mathcal{C}}$. The determinant of the $\mathcal{O}_S$ -endomorphism p_*F

$$\Delta := \det(p_*F) \quad (16)$$

is called the *discriminant*. $\mathcal{D} := \varphi(\mathcal{C})$ is defined by the ideal sheaf $\mathcal{J} = (\Delta)$.

Let us put

$$Der_S(-\log \mathcal{D}) := \{\delta \in Der_S \mid \delta(\Delta) \in (\Delta)\}. \quad (17)$$

which is an $\mathcal{O}_S$ -free module of rank μ , closed under bracket product. Considering the $\mathcal{O}_T$ -homomorphism

$$r : \pi_*Der_S \rightarrow q_*\mathcal{O}_{\mathcal{C}}, \quad (18)$$

we see that $\ker(r) = \pi_* \text{Der}_S(-\log \mathcal{D})$ and we have a direct sum decomposition

$$\pi_* \text{Der}_S = \mathcal{G} \oplus \pi_* \text{Der}_S(-\log \mathcal{D}), \quad (19)$$

and an $\mathcal{O}_T$ -homomorphism w

$$w : \mathcal{G} \rightarrow \pi_* \text{Der}_S(-\log \mathcal{D}), \quad \delta \mapsto w(\delta) = t^0 \delta - t^0 \circ \delta. \quad (20)$$

Let us define the specific vector field $E := w(\delta_0)$ which we call the *Euler vector field*. The discriminant Δ is homogeneous degree μ , with respect to E , i.e.,

$$E\Delta = \mu\Delta. \quad (21)$$

1.3 De Rham cohomology $\mathcal{H}_F^{(0)}$

Let Ω_X^p be the sheaf of germs of holomorphic p -forms on X , $\Omega_{X/T}^p := \Omega_X^p / dq^* \Omega_T^1 \wedge \Omega_X^{p-1}$ and $\Omega_{X/S}^p := \Omega_X^p / d\varphi^* \Omega_S^1 \wedge \Omega_X^{p-1}$ be the sheaves of germs of holomorphic relative p -forms with respect to q or φ respectively. In particular let us define,

$$\Omega_F := \Omega_{X/S}^{n+1}. \quad (22)$$

Note that if we fix the holomorphic $n+1$ -form $dz_0 \wedge \cdots \wedge dz_n$, we have $\Omega_F \simeq \mathcal{O}_C dz_0 \wedge \cdots \wedge dz_n$.

Let us define the $\mathcal{O}_S$ -free modules,

$$\begin{aligned} \mathcal{H}_F^{(0)} &:= \varphi_* \Omega_{X/T}^{n+1} / dF^0 \wedge d(\varphi_* \Omega_{X/T}^{n-1}) \\ \mathcal{H}_F^{(-1)} &:= \varphi_* \Omega_{X/T}^n / dF^0 \wedge (\varphi_* \Omega_{X/T}^{n-1}) + d(\varphi_* \Omega_{X/T}^{n-1}) \\ &= \varphi_* \Omega_{X/S}^n / d(\varphi_* \Omega_{X/S}^{n-1}) \\ \mathcal{H}_F^{(-2)} &:= \ker(d : \varphi_* \Omega_{X/S}^n \rightarrow \varphi_* \Omega_{X/S}^{n+1}) / d(\varphi_* \Omega_{X/S}^{n-1}) \\ &= \mathbb{R}^n \varphi_*(\Omega_{X/S}, d), \end{aligned} \quad (23)$$

where F^0 is a function on X defined by $F = t^0 - F^0$. They are $\mathcal{O}_S$ -free modules of rank μ because of the Milnor's fibration theorem:

Theorem 1.2 (Milnor's fibration theorem). *The restriction*

$$\varphi|_{X-\varphi^{-1}(\mathcal{D})} : X - \varphi^{-1}(\mathcal{D}) \rightarrow S - \mathcal{D} \quad (24)$$

is a locally trivial fibration, whose general fiber $X_s = \varphi^{-1}(s)$ for $s \in S - \mathcal{D}$ is homeomorphic to a bouquet of n -spheres.

Indeed, we see that

$$H^i(X_s, \mathbb{C}) \simeq \begin{cases} \mathbb{C}, & i = 0, \\ 0, & i \neq 0 \text{ and } n, \\ \mathbb{C}^\mu, & i = n. \end{cases} \quad (25)$$

The exterior differentiation $d : \Omega_{X/T}^n \rightarrow \Omega_{X/T}^{n+1}$ induces an $\mathcal{O}_S$ -exact sequence,

$$0 \rightarrow \mathcal{H}_F^{(-2)} \hookrightarrow \mathcal{H}_F^{(-1)} \xrightarrow{d} \varphi_* \Omega_F \rightarrow 0. \quad (26)$$

The exterior product $dF^0 : \Omega_{X/T}^n \rightarrow \Omega_{X/T}^{n+1}$ induces an $\mathcal{O}_S$ -exact sequence,

$$0 \rightarrow \mathcal{H}_F^{(-1)} \xrightarrow{dF^0} \mathcal{H}_F^{(0)} \rightarrow \varphi_* \Omega_F \rightarrow 0. \quad (27)$$

Definition 1.3. The covariant differentiation

$$\nabla : \text{Der}_S \times \mathcal{H}_F^{(-1)} \rightarrow \mathcal{H}_F^{(0)}, \quad (28)$$

defined by

$$\nabla_{\frac{\partial}{\partial t^i}}[\zeta] = (-1)^i [(dt^1 \wedge \cdots \wedge dt^{\mu-1})^{-1} dt^0 \wedge \cdots \wedge \hat{dt}^i \wedge \cdots \wedge dt^{\mu-1} \wedge d\zeta], \quad (29)$$

is called the *Gauss–Manin connection*.

Proposition 1.3. *The Gauss–Manin connection is integrable and*

$$\nabla_{\delta_0} : \mathcal{H}_F^{(-1)} \simeq \mathcal{H}_F^{(0)}, \quad \nabla_{\delta_0}[\zeta] = [d\zeta]. \quad (30)$$

Then the map

$$\nabla_{\delta_0}^{-1} : \mathcal{H}_F^{(0)} \rightarrow \mathcal{H}_F^{(-1)}, \quad [\zeta] \mapsto [dF^0 \wedge d^{-1}\zeta], \quad (31)$$

is well-defined.

Thus we can consider the following decreasing sequence

$$\mathcal{H}_F^{(-k-1)} := \{\omega \in \mathcal{H}_F^{(-k)} : \nabla_{\delta_0} \omega \in \mathcal{H}_F^{(-k)}\}, \quad k \in \mathbb{N} \quad (32)$$

of $\mathcal{O}_S$ -free submodules of $\mathcal{H}_F^{(0)}$.

We get the $\mathcal{O}_S$ -exact sequences,

$$0 \rightarrow \mathcal{H}_F^{(-k-1)} \hookrightarrow \mathcal{H}_F^{(-k)} \xrightarrow{r^{(k)}} \varphi_* \Omega_F \rightarrow 0, \quad k \in \mathbb{N}, \quad (33)$$

and the Gauss–Manin connection ∇ ,

$$\nabla : \text{Der}_S \times \mathcal{H}_F^{(-k-1)} \rightarrow \mathcal{H}_F^{(-k)}, \quad k \in \mathbb{N}. \quad (34)$$

Remark. Since $r^{(k)}(\nabla_\delta \zeta) = r(\delta)r^{(k)}(\nabla_{\delta_0} \zeta)$, we have

$$\nabla : \text{Der}_S(-\log(\mathcal{D})) \times \mathcal{H}_F^{(-k)} \rightarrow \mathcal{H}_F^{(-k)}, \quad k \in \mathbb{N}. \quad (35)$$

This means that the Gauss–Manin connection is regular singular.

1.4 Higher residue pairings $K^{(k)}$ and primitive form $\zeta^{(0)}$

We will consider the direct images $\pi_*\mathcal{H}_F^{(-k)}$. Note that they are not finite $\mathcal{O}_T$ -modules. The pairing $K^{(k)}$ on $\pi_*\mathcal{H}_F^{(0)}$ in the following theorem is called the *higher residue pairings* [S1][S2].

Theorem 1.4. *Let $F(z, t)$ be an LG system over a frame (Z, X, S, T) . Then there exists an infinite sequence of $\mathcal{O}_T$ -bilinear forms,*

$$K^{(k)} : \pi_*\mathcal{H}_F^{(0)} \times \pi_*\mathcal{H}_F^{(0)} \rightarrow \mathcal{O}_T, \quad k \in \mathbb{N} \quad (36)$$

with the following properties:

1. $K^{(k)}$ is symmetric (skew-symmetric) when k is even (odd), respectively.

$$2. \quad K^{(0)}([\phi_1 dz], [\phi_2 dz]) = \text{Res}_{X/T} \left[\begin{array}{c} \phi_1 \phi_2 dz_0 \wedge \cdots \wedge dz_n \\ \frac{\partial F}{\partial z_0} \cdots \frac{\partial F}{\partial z_n} \end{array} \right],$$

for $[\phi_1 dz_0 \wedge \cdots \wedge dz_n], [\phi_2 dz_0 \wedge \cdots \wedge dz_n] \in \pi_*\mathcal{H}_F^{(0)}$.

3. $K^{(k)}(\omega_1, \omega_2) = K^{(k-1)}(\nabla_{\delta_0} \omega_1, \omega_2)$, for $\omega_1 \in \pi_*\mathcal{H}_F^{(-1)}$, $\omega_2 \in \pi_*\mathcal{H}_F^{(0)}$.

4. $K^{(k)}(t^0 \omega_1, \omega_2) - K^{(k)}(\omega_1, t^0 \omega_2) = (n+k)K^{(k-1)}(\omega_1, \omega_2)$,
for $\omega_1, \omega_2 \in \pi_*\mathcal{H}_F^{(0)}$.

5. $\delta K^{(k)}(\omega_1, \omega_2) = K^{(k)}(\nabla_{\delta} \omega_1, \omega_2) + K^{(k)}(\omega_1, \nabla_{\delta} \omega_2)$,
for $\omega_1, \omega_2 \in \pi_*\mathcal{H}_F^{(-1)}$, $\delta \in \mathcal{G}$.

Such a $K^{(k)}$ with the above properties is unique up to constant factor.

Remark. $K^{(0)}$ induces a non-degenerate $\mathcal{O}_T$ -bilinear form J :

$$J : q_*\Omega_F \times q_*\Omega_F \rightarrow \mathcal{O}_T. \quad (37)$$

Definition 1.4. An element $\zeta^{(0)} \in \Gamma(S, \mathcal{H}_F^{(0)})$ is called a *primitive form* if it satisfies the following conditions:

1. Invertibility. $r^{(0)}(\zeta^{(0)}) \in \Gamma(C, \Omega_F)$ is an $\mathcal{O}_C$ -free base of Ω_F .
2. First integrability. $K^{(1)}(\nabla_{\delta} \zeta^{(-1)}, \nabla_{\delta'} \zeta^{(-1)}) = 0$ for $\delta, \delta' \in \mathcal{G}$.
3. Homogeneity. $\nabla_E \zeta^{(0)} = (r-1)\zeta^{(0)}$ for a constant r .
4. $K^{(k)}(\nabla_{\delta} \nabla_{\delta'} \zeta^{(-2)}, \nabla_{\delta''} \zeta^{(-1)}) = 0$ for $k \geq 2$ and $\delta, \delta', \delta'' \in \mathcal{G}$.
5. $K^{(k)}(t^0 \nabla_{\delta} \zeta^{(-1)}, \nabla_{\delta'} \zeta^{(-1)}) = 0$ for $k \geq 2$ and $\delta, \delta' \in \mathcal{G}$.

where $\zeta^{(-k)} := (\nabla_{\delta_0})^{-k} \zeta$.

Theorem 1.5. *If S is small enough, then there exists a primitive form $\zeta^{(0)}$.*

1.5 Canonical coordinate system

Since $r^{(0)}(\zeta^{(0)})$ is invertible, there is an $\mathcal{O}_T$ -isomorphism

$$\mathcal{G} \simeq q_*\Omega_F, \quad \delta \mapsto \delta F|_{\mathcal{C}} \cdot r^{(0)}(\zeta^{(0)}). \quad (38)$$

Then we can define a connection

$$\tilde{\nabla} : \text{Der}_T \times \mathcal{G} \rightarrow \mathcal{G} \quad (39)$$

by the relation

$$\nabla_{\delta} \nabla_{\delta'} \zeta^{(-2)} = \nabla_{\delta \circ \delta'} \zeta^{(-1)} + \nabla_{\tilde{\nabla}_{\delta} \delta'} \zeta^{(-2)}, \quad (40)$$

which depends only on $r^{(0)}(\zeta^{(0)})$.

The integrability of the Gauss–Manin connection ∇ on $\mathcal{H}_F^{(0)}$ follows that

$$\tilde{\nabla}_{\delta}(\delta' \circ \delta'') - \delta' \circ \tilde{\nabla}_{\delta} \delta'' - \tilde{\nabla}_{\delta'}(\delta \circ \delta'') + \delta \circ \tilde{\nabla}_{\delta'} \delta'' = [\delta, \delta'] \circ \delta'', \quad \delta, \delta', \delta'' \in \mathcal{G}, \quad (41)$$

and

$$[\tilde{\nabla}_{\delta}, \tilde{\nabla}_{\delta'}] = \tilde{\nabla}_{[\delta, \delta']}, \quad \delta, \delta' \in \mathcal{G}. \quad (42)$$

By putting $\delta'' = \delta_0$ in (41), we see that this connection is torsion free

$$\tilde{\nabla}_{\delta} \delta' - \tilde{\nabla}_{\delta'} \delta = [\delta, \delta'], \quad \delta, \delta' \in \mathcal{G}. \quad (43)$$

From the fifth property of $K^{(k)}$, we see that this connection is metric with respect to J ,

$$\delta J(\delta', \delta'') = J(\tilde{\nabla}_{\delta} \delta', \delta'') + J(\delta', \tilde{\nabla}_{\delta} \delta''), \quad \delta, \delta', \delta'' \in \mathcal{G}, \quad (44)$$

where we denote $J(\delta' F|_{\mathcal{C}} \cdot r^{(0)}(\zeta^{(0)}), \delta'' F|_{\mathcal{C}} \cdot r^{(0)}(\zeta^{(0)}))$ by $J(\delta, \delta')$ for short.

We can define an $\mathcal{O}_T$ -endomorphism N

$$N : \mathcal{G} \rightarrow \mathcal{G} \quad (45)$$

by the relation

$$t^0 \nabla_{\delta} \zeta^{(-1)} = \nabla_{t^0 \circ \delta} \zeta^{(-1)} + \nabla_{N\delta} \zeta^{(-2)}, \quad \delta \in \mathcal{G}. \quad (46)$$

The commutation relation $[\delta, t^0] = \delta t^0$ follows

$$\tilde{\nabla}_{\delta}(t^0 \circ \delta') + \delta \circ (N + 1)\delta' = N(\delta \circ \delta') + t^0 \circ \tilde{\nabla}_{\delta} \delta' + (\delta t^0)\delta', \quad \delta, \delta' \in \mathcal{G}, \quad (47)$$

and

$$\tilde{\nabla}_\delta(N\delta') = N(\tilde{\nabla}_\delta\delta'), \quad \delta, \delta' \in \mathcal{G}. \quad (48)$$

From the fourth property of $K^{(k)}$, we get

$$J(N\delta, \delta') + J(\delta, N\delta') = (n+1)J(\delta, \delta'), \quad \delta, \delta' \in \mathcal{G}, \quad (49)$$

i.e.

$$N + N^* = (n+1)id, \quad (50)$$

where N^* is the adjoint of N with respect to the metric J .

Since the connection $\tilde{\nabla}$ on $\mathcal{G}$ is torsion free and integrable, there exists a unique coordinate system (called a *flat coordinate system*) $t^0, \dots, t^{\mu-1}$ of S up to linear transformations such that the horizontal vector space of $\tilde{\nabla}$ is given by

$$\ker \tilde{\nabla} = \bigoplus_{i=0}^{\mu-1} \mathbb{C} \frac{\partial}{\partial t^i}. \quad (51)$$

The equation (48) implies that N is a $\mathbb{C}$ -linear endomorphism of $\ker \tilde{\nabla}$.

Definition 1.5. The eigenvalues $\{\alpha_0, \alpha_1, \dots, \alpha_{\mu-1}\}$ of N is called the *exponents*.

We introduce the following *Poincaré polynomial* associated to the LG system F :

$$\chi(F)(y, \bar{y}) := \sum_{i=0}^{\mu-1} (y\bar{y})^{q_i}, \quad (52)$$

where $q_i := \alpha_i - r$, $i = 0, \dots, \mu-1$.

Since we can choose exponents such that $\alpha_0(= r) \leq \dots \leq \alpha_{\mu-1}$ and exponents have the following duality property:

$$\{\alpha_0, \dots, \alpha_{\mu-1}\} = \{n+1-\alpha_0, \dots, n+1-\alpha_{\mu-1}\}, \quad (53)$$

it is obvious from (50) that the duality property of the Poincaré polynomial,

$$\chi(F)(y, \bar{y}) = (y\bar{y})^{\hat{c}_F} \chi(F)(y^{-1}, \bar{y}^{-1}). \quad (54)$$

where $\hat{c}_F := q_{\mu-1} = (n+1) - 2r$.

Let $t^0, \dots, t^{\mu-1}$ be a flat coordinate system. If $N\left(\frac{\partial}{\partial t^i}\right) = \alpha_i \frac{\partial}{\partial t^i}$, then we can define the *degree* of t^i by

$$\deg(t^i) := 1 - q_i, \quad i = 0, \dots, \mu-1. \quad (55)$$

Definition 1.6. A coordinate system $t^0, \dots, t^{\mu-1}$ of S is called a *canonical coordinate system* if it is a flat coordinate system and

$$N\left(\frac{\partial}{\partial t^i}\right) = \alpha_i \frac{\partial}{\partial t^i}, \quad i = 1, \dots, \mu - 1. \quad (56)$$

Remark. In a canonical coordinate system, the Euler vector field $E = w(\delta_0)$ is given by

$$E = \sum_{i=0}^{\mu-1} \deg(t^i) t^i \frac{\partial}{\partial t^i}. \quad (57)$$

Remark. This canonical coordinates coincides with the one introduced by [BCOV].

2 Cohomological Field Theory associated to LG system

2.1 Frobenius structure

In this subsection, we will show that S is a Frobenius manifold. We first review the Frobenius structure [D][Man1].

Definition 2.1. An affine flat structure on the supermanifold M is a sub-sheaf $\mathcal{T}_M^f \subset \mathcal{T}_M$ of linear spaces of pairwise (super)commuting vector fields, such that $\mathcal{T}_M = \mathcal{T}_M^f \otimes \mathcal{O}_M$ (tensor product over the ground field). Sections of $\mathcal{T}_M^f$ is called flat vector field.

A Riemannian metric on M is an even symmetric pairing $\eta : S^2(\mathcal{T}_M) \rightarrow \mathcal{O}_M$, inducing an isomorphism $\eta' : \mathcal{T}_M \rightarrow \mathcal{T}_M^*$. We put $\eta_{ab} := \eta(\partial_a, \partial_b)$.

Definition 2.2. A Riemannian metric $\eta \in \mathcal{T}_M^* \otimes \mathcal{T}_M^*$ is compatible with the structure $\mathcal{T}_M^f$ if $\eta(X, Y)$ is constant for all $X, Y \in \mathcal{T}_M^f$.

Definition 2.3. Let M be a supermanifold. Consider a triple $(\mathcal{T}_M^f, \eta, C)$ consisting of an affine flat structure, a compatible metric and an even symmetric tensor $C : S^3(\mathcal{T}_M) \rightarrow \mathcal{O}_M$.

Define an $\mathcal{O}_M$ -bilinear symmetric multiplication $\circ$ on $\mathcal{T}_M$:

$$\mathcal{T}_M \otimes \mathcal{T}_M \longrightarrow S^2(\mathcal{T}_M) \xrightarrow{C'} \mathcal{T}_M^* \xrightarrow{\eta'} \mathcal{T}_M, \quad (58)$$

where primes denotes a partial dualization.

Then M endowed with this structure is called a *pre-Frobenius* manifold.

A local even function $\Phi \in \mathcal{O}_M$ is called a *Frobenius potential* if

$$C(X, Y, Z) = (XYZ)\Phi, \quad X, Y, Z \in \mathcal{T}_M. \quad (59)$$

Definition 2.4. A pre-Frobenius manifold is called *Frobenius*, if C everywhere locally admits a potential and the multiplication $\circ$ is associative.

Remark. If a Frobenius potential exists, it is unique up to a quadratic polynomial in flat local coordinates.

Remark. In flat local coordinates t^a , then

$$C_{abc} = \partial_a \partial_b \partial_c \Phi, \quad (60)$$

$$\partial_a \circ \partial_b = \sum_c C_{ab}^c \partial_c, \quad (61)$$

where

$$C_{ab}^c := \sum_d C_{abd} \eta^{cd}, \quad (\eta^{ab}) := (\eta_{ab})^{-1}. \quad (62)$$

Definition 2.5. Let (M, η, Φ) be a Frobenius manifold. An even vector field $e \in \mathcal{T}_M$ is called a *identity* if $e \circ X = X$ for all vector fields $X \in \mathcal{T}_M$.

Definition 2.6. An even vector field $E \in \mathcal{T}_M$ is called an *Euler vector field* if it satisfies the following:

$$E(\eta(X, Y)) - \eta([E, X], Y) - \eta(X, [E, Y]) = D\eta(X, Y), \quad X, Y \in \mathcal{T}_M, \quad (63)$$

and

$$[E, X \circ Y] - [E, X] \circ Y - X \circ [E, Y] = d_0 X \circ Y, \quad X, Y \in \mathcal{T}_M. \quad (64)$$

Theorem 2.1. Let $F(z, t)$ be an LG system over a frame (Z, X, S, T) and let us assume that a primitive form $\zeta^{(0)}$ is given. Then S is a Frobenius manifold with an identity δ_0 and an Euler vector field $E = w(\delta_0)$.

Proof. It is clear that a canonical coordinate system t^i , $i = 1, \dots, \mu$ give an affine flat structure. We will show the existence of a compatible metric η and a symmetric tensor C .

Let

$$\eta_F(\delta, \delta') := \sum_{i,j=0}^{\mu-1} f^i g^j J(\delta_i, \delta_j), \quad (65)$$

for $\delta = \sum_{i=0}^{\mu-1} f^i \delta_i$, $\delta' = \sum_{j=0}^{\mu-1} f^j \delta_j \in \text{Der}_S$. Then $\eta_F(\delta_i, \delta_j)$ is constant for all flat vector fields δ_i, δ_j because of the second property of the definition 1.4;

$$\begin{aligned} \delta_k \eta_F(\delta_i, \delta_j) &= \delta_k K^{(0)}(\nabla_{\delta_i} \zeta^{(-1)}, \nabla_{\delta_j} \zeta^{(-1)}) \\ &= K^{(1)}(\nabla_{\delta_k} \nabla_{\delta_i} \zeta^{(-2)}, \nabla_{\delta_j} \zeta^{(-1)}) - K^{(1)}(\nabla_{\delta_i} \zeta^{(-1)}, \nabla_{\delta_k} \nabla_{\delta_j} \zeta^{(-2)}) \\ &= K^{(1)}(\nabla_{\delta_k \circ \delta_i} \zeta^{(-1)}, \nabla_{\delta_j} \zeta^{(-1)}) - K^{(1)}(\nabla_{\delta_i} \zeta^{(-1)}, \nabla_{\delta_k \circ \delta_j} \zeta^{(-1)}) \\ &= 0 \end{aligned} \quad (66)$$

Let us define C by the residual product $\circ$

$$\delta_i \circ \delta_j = \sum_{k=0}^{\mu} C_{ij}^k \delta_k. \quad (67)$$

Since

$$\begin{aligned} \partial_i C_{jkl} &= \delta_i K^{(0)}(\nabla_{\delta_j} \nabla_{\delta_k} \zeta^{(-2)}, \nabla_{\delta_l} \zeta^{(-1)}) \\ &= K^{(0)}(\nabla_{\delta_i} \nabla_{\delta_j} \nabla_{\delta_k} \zeta^{(-2)}, \nabla_{\delta_l} \zeta^{(-1)}) \\ &= K^{(0)}(\nabla_{\delta_j} \nabla_{\delta_i} \nabla_{\delta_k} \zeta^{(-2)}, \nabla_{\delta_l} \zeta^{(-1)}) \\ &= \delta_j K^{(0)}(\nabla_{\delta_i} \nabla_{\delta_k} \zeta^{(-2)}, \nabla_{\delta_l} \zeta^{(-1)}) \\ &= \partial_j C_{ikl}, \end{aligned} \quad (68)$$

there exists a potential Φ_F such that $C_{ijk} = \partial_i \partial_j \partial_k \Phi_F$. The associativity is obvious from the definition of the residual product $\circ$. The rest is clear. $\square$

2.2 Cohomological Field Theory

Let k be a supercommutative $\mathbb{Q}$ -algebra, H a $(\mathbb{Z}_2\text{-graded})$ free k -module of finite rank, $\eta : H \times H \rightarrow k$ an even symmetric non-degenerate pairing on H . We introduce formally the *large phase space* as linear infinite dimensional

formal supermanifold $\oplus_{d \geq 0} H[d]$ with basis $\sigma_d(\mathcal{O}_a)$ and coordinates t_d^a . Put $t^a := t_0^a$, $t := \{t^a\}$.

A *Frobenius potential* on (H, η) is given by an even potential $\Phi \in k[[t]]$, defined up to quadratic terms, whose third derivatives form the structure constants of the supercommutative, associative $k[[t]]$ -algebra $H \otimes k[[t]]$.

Definition 2.7. A triple $M = (H, \eta, \Phi)$ is called a *formal Frobenius manifold* (over k).

Let $\overline{\mathcal{M}}_{0,n}$ be the Deligne–Mumford compactification of the moduli space of genus 0 Riemann surfaces with n -marked points. The pair k -vector space H and maps I_n^M in the following proposition is called the (tree level) *Cohomological Field Theory* [KM1][Man1].

Proposition 2.2. [KM2] *For any formal Frobenius manifold M , there exists a unique sequence of k -linear maps*

$$I_n^M : H^n \rightarrow H^*(\overline{\mathcal{M}}_{0,n}, k), \quad n \geq 3, \quad (69)$$

satisfying the following properties:

1. $\mathcal{S}_n$ -covariance (with respect to the natural action of $\mathcal{S}_n$ on both sides of (69)).
2. *Splitting, or compatibility with restriction to the boundary divisors: for any stable ordered partition $\sigma : \{1, \dots, n\} = S_1 \amalg S_2$, $n_i = |S_i|$, and the respective map*

$$\varphi_\sigma : \overline{\mathcal{M}}_{0,n_1+1} \times \overline{\mathcal{M}}_{0,n_2+1} \rightarrow \overline{\mathcal{D}}(\sigma) \subset \overline{\mathcal{M}}_{0,n} \quad (70)$$

we have

$$\begin{aligned} \varphi_\sigma^*(I_n^M(\mathcal{O}_{a_1} \otimes \dots \otimes \mathcal{O}_{a_n})) \\ = \epsilon(\sigma)(I_{n_1+1}^M \otimes I_{n_2+1}^M)((\bigotimes_{p \in S_1} \mathcal{O}_{a_p}) \otimes \Delta \otimes (\bigotimes_{q \in S_2} \mathcal{O}_{a_q})) \end{aligned} \quad (71)$$

where $\Delta := \sum_{a,b} \mathcal{O}_a \eta^{ab} \mathcal{O}_b$, and $\epsilon(\sigma)$ is the sign of the permutation induced on the odd arguments $\mathcal{O}_{a_1}, \dots, \mathcal{O}_{a_n}$.

3.

$$\int_{\overline{\mathcal{M}}_{0,n}} I_n^M(\mathcal{O}_{a_1} \otimes \dots \otimes \mathcal{O}_{a_n}) = \partial_{a_1} \dots \partial_{a_n} \Phi|_{t=0}, \quad (72)$$

where the integral denotes the value of the top degree term of I_n^M on the fundamental class $\overline{\mathcal{M}}_{0,n}$.

Now we can define the correlators with gravitational descendants $\sigma_d(\mathcal{O}_a)$ by

$$\begin{aligned} & \langle \sigma_{d_1}(\mathcal{O}_{a_1}) \dots \sigma_{d_n}(\mathcal{O}_{a_n}) \rangle \\ &:= \int_{\overline{\mathcal{M}}_{0,n}} I_n^M(\mathcal{O}_{a_1} \otimes \dots \otimes \mathcal{O}_{a_n}) c_1(\mathcal{L}_1)^{d_1} \dots c_1(\mathcal{L}_n)^{d_n}, \end{aligned} \quad (73)$$

where $\mathcal{L}_i$ is the line bundle which has the geometric fiber $T_{x_i}^*C$ at the point $(C; x_1, \dots, x_n)$.

Definition 2.8. A formal function Φ^{grav} on large phase space $\oplus_{d \geq 0} H[d]$ defined by

$$\Phi^{grav} := \sum_{n, (d_1, a_1), \dots, (d_n, a_n)} \epsilon(a) \frac{1}{n!} t_{d_1}^{a_1} \dots t_{d_n}^{a_n} \langle \sigma_{d_1}(\mathcal{O}_{a_1}) \dots \sigma_{d_n}(\mathcal{O}_{a_n}) \rangle \quad (74)$$

is called a *genus 0 free energy*.

Let us introduce μ -dimensional vector spaces,

$$H_F^{(d)} := \bigoplus_{i=0}^{\mu-1} \mathbb{C} \nabla_{\delta_i} \zeta^{(-d-1)} \big|_{t'=0} \quad (75)$$

$$\simeq \left(\pi_* \mathcal{H}_F^{(-d)} \bigg/ \pi_* \mathcal{H}_F^{(-d-1)} \right) \bigg/ m_0 \left(\pi_* \mathcal{H}_F^{(-d)} \bigg/ \pi_* \mathcal{H}_F^{(-d-1)} \right), \quad d \in \mathbb{N}, \quad (76)$$

where $m_0 = (t') := (t^1, \dots, t^{\mu-1})$ is the maximal ideal of $\mathcal{O}_{T,0}$.

Lemma 2.3. Let $F(z, t)$ be an LG system over a frame (Z, X, S, T) and let us assume that a primitive form $\zeta^{(0)}$ is given. Then $M_F = (H_F^{(0)}, \eta_F, \Phi_F)$ is a formal Frobenius manifold over $\mathbb{C}$.

Proof. Since S is a Frobenius manifold, it is clear that M_F is a formal Frobenius manifold over $\mathbb{C}$. □

Remark. The large phase space is given by

$$H_F := \pi_* \mathcal{H}_F^{(0)} \bigg/ m_0 \pi_* \mathcal{H}_F^{(0)} \simeq \bigoplus_{d \geq 0} H_F^{(d)} \quad (77)$$

and

$$\sigma_d(\mathcal{O}_i) = \nabla_{\delta_i} \zeta^{(-d-1)} \big|_{t'=0}, \quad i = 0, \dots, \mu-1, \quad d \geq 0. \quad (78)$$

This is the multi-variable version of a similar formula given by Losev [L]. We can obtain the correlators with the gravitational descendants by using the the higher residues and the above expressions.

Thus we have built up the mathematical definition of the topological Landau–Ginzburg model coupled to gravity at genus 0:

Theorem 2.4. *Let $F(z, t)$ be an LG system over a frame (Z, X, S, T) . Then a primitive form $\zeta^{(0)}$ uniquely defines the genus 0 free energy Φ_F^{grav} such that $\Phi_F^{grav}|_{t_d^0=0, d>0} = \Phi_F$ up to quadratic terms.*

Remark. It is well known that (topological recursion relation at genus 0 [W1][W2])

$$\frac{\partial}{\partial t^i} \frac{\partial}{\partial t^j} \left(\frac{\partial}{\partial t_d^l} \Phi_F^{grav} \right) = \sum_{k=0}^{\mu-1} C_{ij}^k \frac{\partial}{\partial t^k} \frac{\partial}{\partial t^0} \left(\frac{\partial}{\partial t_d^l} \Phi_F^{grav} \right). \quad (79)$$

If we restrict the above equation to the small phase space $H_F^{(0)}$ ($t_d^i = 0, d > 0$), it becomes the ordinary Gauss–Manin differential equations for period integrals expressed in a flat coordinate system [S1].

3 Primitive Forms and Mirror Symmetry

3.1 Period of Primitive Forms

In this subsection, we will review the period mapping of primitive forms [S1].

Definition 3.1. For any $\kappa \in \mathbb{C}$, we define the D_S -module

$$\mathcal{M}^{(\kappa)} := D_S / I_S, \quad (80)$$

where

$$I_S := \sum_{\delta, \delta' \in \mathcal{G}} D_S P(\delta, \delta') + \sum_{\delta \in \mathcal{G}} D_S Q_\kappa(\delta), \quad (81)$$

$$P(\delta, \delta') := \delta \delta' - \delta \circ \delta' \delta_0 - \nabla_\delta \delta', \quad \delta, \delta' \in \mathcal{G}, \quad (82)$$

$$Q_\kappa(\delta) := w(\delta) \delta_0 - (N - \kappa - 1) \delta, \quad \delta \in \mathcal{G}. \quad (83)$$

Lemma 3.1. *The module $\mathcal{M}^{(\kappa)}$ for $\kappa \in \mathbb{C}$ is a simple holonomic system.*

For $[P] \in \mathcal{M}^{(\kappa+1)}$, we can define a left D_S -homomorphism

$$\mathcal{M}^{(\kappa+1)} \rightarrow \mathcal{M}^{(\kappa)}, \quad [P] \mapsto [P\delta_0], \quad (84)$$

since we have

$$P(\delta, \delta')\delta_0 = \delta_0 P(\delta, \delta'), \quad (85)$$

and

$$Q_{\kappa+1}(\delta)\delta_0 = \delta_0 Q_{\kappa}(\delta). \quad (86)$$

Let l be the corank of the $N - \kappa - 1$ on the μ -dimensional horizontal vector space of ∇ on $\mathcal{G}$ and let $\eta_1, \dots, \eta_l$ and $\xi_1, \dots, \xi_l$ be the basis of $\ker(N - \kappa - 1)$ and $\text{coker}(N - \kappa - 1)$, respectively. Then we have an D_S -exact sequence

$$0 \rightarrow \mathcal{O}_S[E_{\kappa+1}] \oplus \bigoplus_{i=1}^l \mathcal{O}_S[w(\eta_i)] \rightarrow \mathcal{M}^{(\kappa+1)} \rightarrow \mathcal{M}^{(\kappa)} \rightarrow \mathcal{O}_S \oplus \bigoplus_{i=1}^l \mathcal{O}_S[\xi_i] \rightarrow 0, \quad (87)$$

for $r \neq \kappa + 1$, and

$$0 \rightarrow \bigoplus_{i=1}^l \mathcal{O}_S[w(\eta_i)] \rightarrow \mathcal{M}^{(\kappa+1)} \rightarrow \mathcal{M}^{(\kappa)} \rightarrow \mathcal{O}_S \oplus \bigoplus_{i=2}^l \mathcal{O}_S[\xi_i] \rightarrow 0, \quad (88)$$

for $r = \kappa + 1$ and $\xi_1 = \delta_0$, where $E_{\kappa+1} = E - (r - \kappa - 1)$.

Let us define

$$\text{Sol}(\mathcal{M}^{(\kappa)}) := \text{Hom}_{D_S}(\mathcal{M}^{(\kappa)}, \mathcal{O}_S), \quad \kappa \in \mathbb{C}. \quad (89)$$

Remark. Since $\mathcal{M}^{(\kappa)}$ has the generator 1, a solution $u \in \text{Sol}(\mathcal{M}^{(\kappa)})$ is identified with $u(1) \in \mathcal{O}_S$.

Then we have the following lemma.

Lemma 3.2. *The kernel of the morphism,*

$$\text{Sol}(\mathcal{M}^{(\kappa+1)}) \xrightarrow{\delta_0} \text{Sol}(\mathcal{M}^{(\kappa)}), \quad (90)$$

is spanned by canonical coordinate of exponent $\kappa + 1$ (or degree $r - \kappa$) and by a constant function 1_S .

Since $Sol(\mathcal{M}^{(\kappa)})$ always contains the constant function, we have

$$dSol(\mathcal{M}^{(\kappa)}) \simeq Sol(\mathcal{M}^{(\kappa)}) / \mathbb{C}, \quad (91)$$

where $dSol(\mathcal{M}^{(\kappa)})$ is the image of $Sol(\mathcal{M}^{(\kappa)})$ in Ω_S^1 under d .

Let $\gamma^0(s), \dots, \gamma^{\mu-1}(s) \in H_n(X_s, \mathbb{Z})$ be a $\mathbb{Z}$ -basis of a horizontal family of homology defined on a simply connected domain of a covering space of $S - \mathcal{D}$. We see that $\int_{\gamma(s)} \zeta^{(k-1)} \in Sol(\mathcal{M}^{(k)})$ for $k \in \mathbb{Z}$. In particular, for $k \leq 0$ the constant function 1_S and the periods $\int_{\gamma^i(s)} \zeta^{(k-1)}$, $i = 0, \dots, \mu - 1$ form a $\mathbb{C}$ -basis of $Sol(\mathcal{M}^{(k)})$.

So there exist natural isomorphisms of local systems,

$$\dots \xrightarrow{\delta_0} dSol(\mathcal{M}^{(-2)})|_{S-\mathcal{D}} \xrightarrow{\delta_0} dSol(\mathcal{M}^{(-1)})|_{S-\mathcal{D}} \quad (92)$$

$$\xrightarrow{\delta_0} dSol(\mathcal{M}^{(0)})|_{S-\mathcal{D}} \simeq \bigcup_{s \in S-\mathcal{D}} H_n(X_s, \mathbb{C}), \quad (93)$$

Proposition 3.3. *There exists a non-degenerate $\mathbb{C}$ -bilinear form,*

$$I_\kappa : dSol(\mathcal{M}^{(\kappa)})|_{S-\mathcal{D}} \times dSol(\mathcal{M}^{(n-\kappa)})|_{S-\mathcal{D}} \rightarrow \mathbb{C}|_{S-\mathcal{D}}, \quad (94)$$

defined by

$$I_\kappa := \sum_{i=1}^{\mu} \delta_i \otimes w(\delta^{i*}) \in \mathcal{M}^{(\kappa)} \otimes \mathcal{M}^{(n-\kappa)}, \quad \kappa \in \mathbb{C} \quad (95)$$

where $\delta_1, \dots, \delta_\mu$ and $\delta^{1*}, \dots, \delta^{\mu*}$ are $\mathcal{O}_T$ -basis of $\mathcal{G}$ and the dual basis with respect to J , respectively.

From this proposition, we have natural isomorphisms of local systems,

$$\bigcup_{s \in S-\mathcal{D}} H^n(X_s, \mathbb{C}) \simeq dSol(\mathcal{M}^{(n)})|_{S-\mathcal{D}} \xrightarrow{\delta_0} dSol(\mathcal{M}^{(n+1)})|_{S-\mathcal{D}} \quad (96)$$

$$\xrightarrow{\delta_0} dSol(\mathcal{M}^{(n+2)})|_{S-\mathcal{D}} \xrightarrow{\delta_0} \dots \quad (97)$$

Theorem 3.4. *Let n be even. For $\gamma, \gamma' \in H_n(X_s, \mathbb{Z})$, the intersection number of these cycles is given by,*

$$\langle \gamma, \gamma' \rangle := (2\pi\sqrt{-1})^{-n} I_0 \left(d \int_{\gamma} \zeta^{(-1)}, d \int_{\gamma'} \zeta^{(n-1)} \right). \quad (98)$$

Corollary. *Let n be even. Then*

$$(2\pi\sqrt{-1})^{-n} \delta_0^n : dSol(\mathcal{M}^{(0)})|_{S-\mathcal{D}} \rightarrow dSol(\mathcal{M}^{(n)})|_{S-\mathcal{D}} \quad (99)$$

coincide with the linear mapping defined by the intersection form.

3.2 Landau–Ginzburg Orbifolds and Calabi–Yau Varieties

Let f be a quasi-homogeneous polynomial of z_i with an isolated singularity at base point 0 such that

$$f(\lambda^{w_0} z_0, \dots, \lambda^{w_n} z_n) = \lambda f(z_0, \dots, z_n). \quad (100)$$

Let us consider the LG system F associated to f . Then we define the affine variety

$$X_1 := \{z \in \mathbb{C}^{n+1} | f(z_0, \dots, z_n) + 1 = 0\}, \quad (101)$$

which is the Milnor fiber of φ at $(1, 0, \dots, 0) \in S$. We can compactify X_1 to $\overline{X_1}$ so that the boundary $V := \overline{X_1} \setminus X_1$ is given by

$$V = \{[z] \in \mathbb{P}(w_0, \dots, w_n) | f([z]) = 0\}. \quad (102)$$

We have two isomorphisms of $\mathbb{C}$ -vector spaces

$$H^n(X_1, \mathbb{C}) \simeq \mathbb{C}[z_0, \dots, z_n] \bigg/ \left(\frac{\partial f}{\partial z_0}, \dots, \frac{\partial f}{\partial z_n} \right) \quad (103)$$

and

$$PH^{n-1}(V, \mathbb{C}) \simeq \mathbb{C}[z_0, \dots, z_n] \bigg/ \left(\frac{\partial f}{\partial z_0}, \dots, \frac{\partial f}{\partial z_n} \right) \bigg|_{int}, \quad (104)$$

where we denote by $PH^{n-1}(V, \mathbb{C})$ the primitive part of the cohomology and $|_{int}$ means the restriction to the integral exponents sector (exponents of the monomial $z_0^{a_0} \dots z_n^{a_n}$ is given by $\sum_{i=0}^n (a_i + 1)w_i$).

Remark. Note that the integral exponents sector corresponds to the *untwisted sector* of the LG orbifold theory $f//G$, where G is the discrete group acting on z_i as $z_i \rightarrow \exp(2\pi\sqrt{-1}w_i)z_i$ [IV][KY].

Lemma 3.5. *Let F be an LG system over a frame (Z, X, S, T) and let us assume that a primitive form $\zeta^{(0)}$ is given. Then $(S', \eta_F|_{S'}, \Phi_F|_{S'})$ is a Frobenius manifold, where*

$$S' := \{t \in S | t^i = 0 \text{ for } N(\frac{\partial}{\partial t^i}) \notin \mathbb{Z}\}. \quad (105)$$

Let us assume that $r = \sum_{i=0}^n w_i = 1$ and the hypersurface V in weighted projective space $\mathbb{P}(w_0, \dots, w_n)$ is a smooth Calabi–Yau manifold. Then the following conjecture is quite natural from the viewpoint of the Landau–Ginzburg orbifold theory and the isomorphism (104).

Conjecture 3.1. Let F be an LG system over a frame (Z, X, S, T) and let us assume that a primitive form $\zeta^{(0)}$ is given. Then $(S', \eta_F|_{S'}, \Phi_F|_{S'})$ should be a Frobenius submanifold of the one constructed by Barannikov–Kontsevich for a Calabi–Yau manifold V .

Then we can formulate the mirror conjecture as follows.

Conjecture 3.2. There should exist a Calabi–Yau manifold V^* such that S' is a Frobenius submanifold of the one constructed by Gromov–Witten theory.

Indeed, if $\dim V = 1$, these conjectures are true since the potential is trivial.

Remark. Since the kernel of the map $(2\pi\sqrt{-1})^{-n}\delta_0^n$ (99) are spanned by the canonical coordinates with integral exponents, there exist the horizontal family of homology $\gamma^0(t), \gamma^i(t) \in H_n(X_t, \mathbb{Z})$ such that

$$1 = \int_{\gamma^0(t)} \zeta^{(0)}, \quad (106)$$

and

$$t^i = \int_{\gamma^i(t)} \zeta^{(0)}, \quad \text{for } \deg(t^i) = 0. \quad (107)$$

Recall that, when we calculate the ‘special coordinates’ in the context of mirror symmetry of Calabi–Yau manifolds, we first take the holomorphic form Ω and calculate the period integrals by Picard–Fuchs or GKZ equations. We choose the special solution $\int_{\Gamma^0} \Omega$ which is single-valued near the *large radius limit* or *maximally unipotent monodromy point* and the solutions $\int_{\Gamma^i} \Omega$ which have simple logarithmic singularity at the maximally unipotent monodromy point. Then we define the normalized holomorphic form by

$$\frac{\Omega}{\int_{\Gamma^0} \Omega}, \quad (108)$$

and define the ‘special coordinates’ as

$$\tau^i = \frac{\int_{\Gamma^i} \Omega}{\int_{\Gamma^0} \Omega}. \quad (109)$$

In [BCOV], it is proved that the canonical coordinates at the maximal unipotent monodromy point coincides with the ‘special coordinates’. So we can conclude that to choose the holomorphic form $\frac{\Omega}{\int_{\Gamma^0} \Omega}$ is equivalent to choose the primitive form $\zeta^{(0)}$ at this point.

3.3 Mirror Symmetry for $\mathbb{CP}^1$

In this subsection, we construct the mirror partner of $\mathbb{CP}^1$ in terms of the primitive form.

In [Bat], Batyrev showed that the quantum cohomology ring of $\mathbb{CP}^1$ is given by the Jacobian ring

$$QH_q^*(\mathbb{CP}^1, \mathbb{C}) \simeq \mathbb{C}[z, z^{-1}] \Big/ \left(z \frac{\partial f}{\partial z} \right), \quad (110)$$

where

$$f := z + qz^{-1}, \quad 0 < |q| < 1. \quad (111)$$

This fact implies that it may be possible to construct a primitive form associated to the holomorphic function

$$f : \mathbb{C}^* \rightarrow \mathbb{C}. \quad (112)$$

Let (z, t^1) , (t^0, t^1) and (t^1) be the coordinate of $X' := \mathbb{C}^* \times \mathbb{C}$, $S' := \mathbb{C}^2$ and $T' := \mathbb{C}$, respectively. Then we define the function on $Z' := X' \times_{T'} S'$

$$F(z, t) := t^0 + z + q \exp(t^1) z^{-1}, \quad (113)$$

where q is a parameter of a punctured unit disk $0 < |q| < 1$. Then we can define a frame

$$\begin{array}{ccc} Z' & \xrightarrow{\hat{\pi}'} & X' \\ p' \downarrow & & \downarrow q' \\ S' & \xrightarrow{\pi'} & T', \end{array} \quad (114)$$

the map $\varphi' : X' \rightarrow S'$, the critical set $C \subset X'$ and the discriminant $D = \varphi'(C)$.

In the previous sections, we have assumed that f is a quasi-homogeneous polynomial of z_i with an isolated singularity at base point $0 \in \mathbb{C}^{n+1}$. In such case, the maps π and q are Stein, which are trivial fibrations with contractible fibers and the map φ is Stein. This condition is sufficient for the existence of higher residue pairings $K^{(k)}$.

The map q' and φ' do not satisfy the above conditions, but it is possible to replace a frame (Z', X', S', T') by (Z, X, S, T) such that this new frame satisfies the conditions. Indeed, we can take a new frame (Z, X, S, T) as follows.

$$T := \{(t^1) \in T' : \|(t^1) - (0)\| < \epsilon\}, \quad (115)$$

$$S := \{(t^0, t^1) \in S' : \|(t^0, t^1) - (2q^{\frac{1}{2}}, 0)\| < \delta_1\} \cup \{(t^0, t^1) \in S' : \|(t^0, t^1) - (-2q^{\frac{1}{2}}, 0)\| < \delta_2\} \cap \pi'^{-1}(T), \quad (116)$$

$$X := \{(z, t^1) \in X' : \|(z, t^1) - (q^{\frac{1}{2}}, 0)\| < r_1\} \cup \{(z, t^1) \in X' : \|(z, t^1) - (q^{\frac{1}{2}}, 0)\| < r_2\} \cap \varphi'^{-1}(S), \quad (117)$$

$$Z := X \times_T S, \quad (118)$$

where ϵ , δ_i and r_i , ($i = 1, 2$) are small real number with $1 \gg r_i \gg \delta_i \gg \epsilon > 0$ ($i=1,2$) and $\|\cdot\|$ is some Euclidean distances. The maps are given by restriction .

We see easily that the correspondence

$$T(S)_0 \simeq \mathbb{C} \frac{\partial}{\partial t^0} \oplus \mathbb{C} \frac{\partial}{\partial t^1} \rightarrow \mathbb{C}[z, z^{-1}] \left/ \left(z \frac{\partial f}{\partial z} \right) \right., \quad \delta \mapsto \hat{\delta} F|_{p^{-1}(0)}, \quad (119)$$

is bijection, so we may call F an LG system.

Theorem 3.6. *There exist higher residue pairing $K^{(k)}$ such that*

$$\begin{aligned} K^{(0)}([\phi_1 \frac{dz}{z}], [\phi_2 \frac{dz}{z}]) &= \sum_{z \frac{\partial F}{\partial z} = 0} \frac{\phi_1 \phi_2}{z \frac{\partial}{\partial z} (z \frac{\partial F}{\partial z})} \\ &= \frac{1}{2\pi\sqrt{-1}} \int_{\gamma} \frac{\phi_1 \phi_2}{z \frac{\partial F}{\partial z}} \frac{dz}{z}, \end{aligned} \quad (120)$$

for $[\phi_1 dz/z], [\phi_2 dz/z] \in \pi_* \mathcal{H}_F^{(0)}$ and γ is a homology cycle of $X \setminus \{z \frac{\partial F}{\partial z} = 0\}$, defined by a general fiber of the mapping

$$X \rightarrow \mathbb{R}^+ \times T, \quad (z, t^1) \mapsto (|z \frac{\partial F}{\partial z}|, t^1). \quad (121)$$

Theorem 3.7. *The primitive form for $F(z, t) = t^0 + z + q \exp(t^1) z^{-1}$ is*

$$\zeta^{(0)} = \frac{dz}{z}. \quad (122)$$

and the minimal exponent is $r = 0$, the Euler vector field is

$$E = t^0 \frac{\partial}{\partial t^0} + 2 \frac{\partial}{\partial t^1}, \quad (123)$$

and the canonical coordinates are given by (t^0, t^1) .

Proof. We will show that $[\frac{dz}{z}]$ satisfy the five conditions of Definition 1.4.

1. It is obvious.
2. Since $K^{(1)}$ is skew symmetric, we only have to show that

$$K^{(1)}(\nabla_{\delta_0}\zeta^{(-1)}, \nabla_{\delta_1}\zeta^{(-1)}) = 0. \quad (124)$$

This follows from

$$\begin{aligned} K^{(1)}(\nabla_{\delta_0}\zeta^{(-1)}, \nabla_{\delta_1}\zeta^{(-1)}) &= \frac{1}{2\pi\sqrt{-1}} \cdot \left(-\frac{1}{2}\right) \int_{\gamma} \frac{q \exp(t^1) z^{-1} dz}{(z \frac{\partial F}{\partial z})^2} \frac{dz}{z} \\ &= 0. \end{aligned} \quad (125)$$

3. The Euler vector field E is given by

$$\begin{aligned} E &= t^0 \frac{\partial}{\partial t^0} - t^0 \circ \frac{\partial}{\partial t^0} \\ &= t^0 \frac{\partial}{\partial t^0} + 2 \frac{\partial}{\partial t^1}. \end{aligned} \quad (126)$$

So we have

$$\begin{aligned} \nabla_E \left[\frac{dz}{z} \right] &= t^0 \nabla_{\delta_0} \left[\frac{dz}{z} \right] + 2 \nabla_{\delta_1} \left[\frac{dz}{z} \right] \\ &= -\frac{(z + q \exp(t^1) z^{-1})^2 dz}{(z - q \exp(t^1) z^{-1})^2} \frac{dz}{z} + 2 \frac{2q \exp(t^1)}{(z - q \exp(t^1) z^{-1})^2} \frac{dz}{z} \\ &= -\frac{dz}{z}. \end{aligned} \quad (127)$$

This shows that $r = 0$.

4. Since

$$\frac{\partial}{\partial t^1} \circ \frac{\partial}{\partial t^1} = q \exp(t^1) \frac{\partial}{\partial t^0} \quad (128)$$

and

$$\begin{aligned} z \frac{\partial}{\partial z} \left(\frac{q \exp(t^1) z^{-1} \cdot q \exp(t^1) z^{-1} - q \exp(t^1)}{z \frac{\partial F}{\partial z}} \right) &= q \exp(t^1) z^{-1} \\ &= \frac{\partial}{\partial t^1} \left(\frac{\partial F}{\partial t^1} \right), \end{aligned} \quad (129)$$

we have

$$\begin{aligned}
& \nabla_{\delta_1} \nabla_{\delta_1} \nabla_{\delta_0}^{-1} \left[\frac{dz}{z} \right] - \nabla_{\delta_1 \circ \delta_1} \left[\frac{dz}{z} \right] \\
&= \nabla_{\delta_0} \left(q \exp(t^1) z^{-1} \cdot q \exp(t^1) z^{-1} - q \exp(t^1) \frac{dz}{z} \right) + \frac{\partial}{\partial t^1} \left(\frac{\partial F}{\partial t^1} \right) \frac{dz}{z} \\
&= -z \frac{\partial}{\partial z} \left(\frac{q \exp(t^1) z^{-1} \cdot q \exp(t^1) z^{-1} - q \exp(t^1)}{z \frac{\partial F}{\partial z}} \right) \frac{dz}{z} + \frac{\partial}{\partial t^1} \left(\frac{\partial F}{\partial t^1} \right) \frac{dz}{z} \\
&= 0.
\end{aligned} \tag{130}$$

It is clear that

$$\nabla_{\delta_0} \nabla_{\delta_1} \nabla_{\delta_0}^{-1} \left[\frac{dz}{z} \right] - \nabla_{\delta_0 \circ \delta_1} \left[\frac{dz}{z} \right] = 0 \tag{131}$$

and

$$\nabla_{\delta_0} \nabla_{\delta_0} \nabla_{\delta_0}^{-1} \left[\frac{dz}{z} \right] - \nabla_{\delta_0 \circ \delta_0} \left[\frac{dz}{z} \right] = 0. \tag{132}$$

5. Since we have proved that $K^{(1)}(\nabla_{\delta} \zeta^{(-1)}, \nabla_{\delta'} \zeta^{(-1)}) = 0$, the condition becomes the following form;

$$t^0 \nabla_{\delta} \left[\frac{dz}{z} \right] = \nabla_{t^0 \circ \delta} \left[\frac{dz}{z} \right] + \nabla_{(N-1)\delta} \nabla_{\delta_0}^{-1} \left[\frac{dz}{z} \right], \quad \delta \in \mathcal{G}. \tag{133}$$

We have

$$t^0 \circ \frac{\partial}{\partial t^1} = -2q \exp(t^1) \frac{\partial}{\partial t^0}. \tag{134}$$

Since

$$t^0 \nabla_{\delta_1} \left[\frac{dz}{z} \right] = - \frac{2q \exp(t^1) (z + q \exp(t^1) z^{-1})^2}{(z - q \exp(t^1) z^{-1})^2} \frac{dz}{z} \tag{135}$$

and

$$2q \exp(t^1) \nabla_{\delta_0} \left[\frac{dz}{z} \right] = + \frac{2q \exp(t^1) (z + q \exp(t^1) z^{-1})^2}{(z - q \exp(t^1) z^{-1})^2} \frac{dz}{z}, \tag{136}$$

we have

$$t^0 \nabla_{\delta_1} \left[\frac{dz}{z} \right] = \nabla_{t^0 \circ \delta_1} \left[\frac{dz}{z} \right]. \tag{137}$$

Now, it is clear that the (t^0, t^1) is a canonical coordinate system. $\square$

We can easily show that the potential Φ_F is given by

$$\Phi_F = (t^0)^2 t^1 + q \exp(t^1) \quad (138)$$

up to quadratic terms. So we have constructed the mirror partner of $\mathbb{CP}^1$.

Theorem 3.8. *Let $f(z) = z + qz^{-1}$ be a holomorphic function on $\mathbb{C}^*$. Then the pair $(\mathbb{C}^*, f)$ is a mirror manifold of $\mathbb{CP}^1$. In other words, we have the equation*

$$\Phi_{\mathbb{CP}^1}^{st}(t) = \Phi_F^{grav}(\tilde{t}), \quad (139)$$

where $\Phi_{\mathbb{CP}^1}^{st}$ is the (large phase space) stable genus 0 generating function for Gromov–Witten Invariants

$$\Phi_{\mathbb{CP}^1}^{st}(t) := \sum_{n \geq 3, (a_i, d_i)} \frac{1}{n!} t_{d_1}^{a_1} \dots t_{d_n}^{a_n} \sum_{\beta \in \mathbb{Z}_{\geq 0}} q^\beta \langle \sigma_{d_1}(\mathcal{O}_{a_1}) \dots \sigma_{d_n}(\mathcal{O}_{a_n}) \rangle_{0, \beta[\mathbb{CP}^1]} \quad (140)$$

and

$$\tilde{t}_d^i := t_d^i + \sum_{(j, e)} t_e^j \sum_{\beta \in \mathbb{Z}_{\geq 0}} q^\beta \langle \sigma_{e-d-1}(\mathcal{O}_j) \mathcal{O}^i \rangle_{0, \beta[\mathbb{CP}^1]}. \quad (141)$$

The quantum cohomology ring of a toric Fano manifold $\mathbb{P}_\Sigma$ associated to the n -dimensional fan Σ is given by the Jacobian ring [Bat]

$$QH_\varphi^*(\mathbb{P}_\Sigma, \mathbb{C}) \simeq \mathbb{C}[z_1^\pm, \dots, z_n^\pm] \Big/ (z_1 \frac{\partial f}{\partial z_1}, \dots, z_n \frac{\partial f}{\partial z_n}), \quad (142)$$

of the Laurent polynomial

$$f_\varphi(z) = \sum_{i=1}^d \exp(\varphi(v_i))^{-1} z^{v_i}, \quad (143)$$

where $v_1, \dots, v_d$ is the set of all generators of 1-dimensional cones in Σ and φ is an element of the complexified Kähler cone of $\mathbb{P}_\Sigma$.

It may be possible to construct a primitive form in these cases, but we are confronted by two difficulties which do not exist in $\mathbb{CP}^1$ case. The first is to find a frame (Z, X, S, T) and the second is to find a (good) LG system $F(z, t)$. However, from the results [EHX2], we can imagine that

Conjecture 3.3. There should exist a primitive form for f which is given by $\zeta^{(0)} = [\frac{dz_1}{z_1} \wedge \dots \wedge \frac{dz_n}{z_n}]$. Then the pair $((\mathbb{C}^*)^n, f)$ is a mirror manifold of a toric Fano manifold $\mathbb{P}_\Sigma$.

It will be important to use the notion of primitive forms if one consider the more general mirror problems. For example, ADE-type theory does not correspond to the usual manifolds, but there may exist a ‘mirror symmetry’ [KY][T] and a structure of quantum cohomology such as Virasoro conditions [EHX1].

References

- [Bat] V. V. Batyrev, Journées de Géométrie Algébrique d'Orsay, Astérisque **218** (1993), 9.
- [BCOV] M. Bershadsky, S. Cecotti, H. Ooguri, C. Vafa, Comm. Math. Phys. **165** (1994), 311.
- [Beh] K. Behrend, Inv. Math. **127** (1997) 601.
- [BF] K. Behrend, B. Fantechi. Inv. Math. **128** (1997) 45.
- [BM] K. Behrend, Yu. I. Manin, Duke Math. J., 85 (1996) 1.
- [BK] S. Barannikov, M. Kontsevich, *Frobenius manifolds and formality of Lie algebras of polyvector field*, alg-geom/9710032.
- [D] B. Dubrovin, in Springer LNM 1620.
- [EHX1] T. Eguchi, K. Hori, C. S. Xiong, Phys. Lett. **B402** (1997) 71.
- [EHX2] T. Eguchi, K. Hori, C. S. Xiong, Int. J. Mod. Phys. **A12** (1997) 1743.
- [EX] T. Eguchi, C. S. Xiong, *Quantum Cohomology at Higher Genus: Topological Recursion Relations and Virasoro Conditions*, hep-th/9801010.
- [G1] E. Getzler, J. Amer. Math. Soc. **10** (1997), 973.
- [G2] E. Getzler, *Topological Recursion Relations in Genus 2*, math/9801003.
- [GY] B. Greene, S. -T. Yau, ed., *Mirror Symmetry II*, AMS/IP, 1997.
- [IV] K. Intriligator, C. Vafa, Nucl. Phys. **B339** (1990) 95.
- [IY] K. Ito, S. -K. Yang, *Flat Coordinates, Topological Landau Ginzburg-Models and the Seiberg-Witten Period Integrals*, hep-th/9708017.
- [KM1] M. Kontsevich, Yu. I. Manin, in [GY].
- [KM2] M. Kontsevich, Yu. I. Manin, *Relations between the correlators of the topological sigma-model coupled to gravity*, alg-geom/9708024.

- [KY] T. Kawai, S.-K. Yang, Prog. Theor. Phys. Suppl. No.**118** (1995) 277.
- [LVW] W. Lerch, C. Vafa, N. P. Warner, Nucl. Phys. **B324** (1989) 427.
- [L] A. Losev, Theoret. and Math. Phys. **100** (1994) 879.
- [LT1] J. Li, G. Tian, *Virtual moduli cycles and Gromov-Witten invariants of algebraic varieties*, Preprint, 1996.
- [LT2] J. Li, G. Tian, *Virtual moduli cycles and Gromov-Witten invariants of general symplectic manifolds*, alg-geom/9608032.
- [LT3] J. Li, G. Tian, *Comparison of the algebraic and the symplectic Gromov-Witten invariants*, alg-geom/9712035.
- [Mar] E. Martinec, in V. Knizhnik Memorial Volume, World Scientific, Singapore 1990.
- [Man1] Yu. I. Manin, *Frobenius manifolds, Quantum cohomology, and Moduli spaces (Chapters I,II,III)*, MPI preprint 96-113.
- [Man2] Yu. I. Manin, *Three constructions of Frobenius manifolds*, math/9801006.
- [RT1] Y. Ruan, G. Tian, Journ. Diff. Geom. **42** (1995) 259.
- [RT2] Y. Ruan, G. Tian, *Higher Genus symplectic Invariants and Sigma model Coupled With Gravity*, alg-geom/9601005.
- [S1] K. Saito, Publ. RIMS, Kyoto University **19** (1983) 1231.
- [S2] K. Saito, J. Fac. Sci. Univ. Tokyo Sect. IA **28** (1982) 775.
- [T] A. Takahashi, *K. Saito's Duality for Regular Weight Systems and Duality for Orbifoldized Poincare Polynomials*, math/9801050.
- [V] C. Vafa, in [Y].
- [VW] C. Vafa, N. P. Warner, Phys. Lett. **218B** (1989) 51.
- [W1] E. Witten, Nucl. Phys. **B340** (1989) 281.
- [W2] E. Witten, Surveys in Diff. Geom. **1** (1991) 243.
- [W3] E. Witten, in [Y].
- [Y] S. -T. Yau, ed., *Essays on Mirror Manifolds*, International Press, Hong Kong, 1992.